



Original Article

The regulation of Oct4 in human gingival fibroblasts stimulated by cyclosporine A: Preliminary observations

Cheng-Chia Yu ^{a,b,1}, Chia-Ming Liu ^{b,c,1}, Tai-Chen Lin ^b,
Ni-Yu Su ^b, Li-Chiu Yang ^b, Yu-Chao Chang ^{b,c*}



^a Institute of Oral Sciences, Chung Shan Medical University, Taichung, Taiwan

^b School of Dentistry, Chung Shan Medical University, Taichung, Taiwan

^c Department of Dentistry, Chung Shan Medical University Hospital, Taichung, Taiwan

Received 23 July 2019; Final revision received 7 October 2019

Available online 27 December 2019

KEYWORDS

Cyclosporine A;
Human gingival
fibroblasts;
Embryonic stem cell
marker;
Oct4;
Nanog

Abstract *Background/purpose:* Oct4, a key transcription factor, could reprogram human somatic fibroblasts into embryonic stem cell-like pluripotent cells. The exact mechanism of cyclosporine A (CsA)-induced gingival overgrowth is still unclear. The aim of this study was to investigate the effects of CsA on the expression of Oct4 in cultured human gingival fibroblasts (HGFs) *in vitro*.

Materials and methods: The effects of CsA on HGFs were used to elucidate whether Oct4 expression could be induced by CsA by using quantitative real-time reverse transcription-polymerase chain reaction and western blot. Cell growth in CsA-treated HGFs with Oct4 lentiviral-mediated shRNAi knockdown was evaluated by tetrazolium bromide reduction assay.

Results: CsA was found to upregulate Oct4 transcript in a dose-dependent manner ($p < 0.05$). CsA also dose-dependently increased Oct4 protein expression ($p < 0.05$). The lentivirus expressing sh-Oct4 successfully prevented the CsA-induced Oct4 mRNA and protein in HGFs ($p < 0.05$). However, knockdown of Oct4 was insufficient to inhibit CsA-stimulated cell growth in HGFs. Furthermore, double knockdown with pluripotency-associated transcription factor Nanog showed that the down-regulation of Oct4/Nanog by lentiviral infection significantly inhibited CsA-stimulated cell growth ($p < 0.05$).

Conclusion: Taken together, CsA was first found to upregulate Oct4 mRNA and protein expression in HGFs. The silencing Oct4 could not suppress cell growth unless Nanog was repressed simultaneously.

© 2019 Association for Dental Sciences of the Republic of China. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author. School of Dentistry, Chung Shan Medical University, 110, Sec. 1, Chien-Kuo N. Rd., Taichung, Taiwan. Fax: +886 4 24759065.

E-mail address: cyc@csmu.edu.tw (Y.-C. Chang).

¹ These authors contributed equally to this work.

Introduction

Cyclosporine A (CsA), a hydrophobic fungal metabolite from *Tolypocladium inflatum*, is an immune-suppressant widely used to prevent transplant rejection and manage several autoimmune conditions.¹ Like the calcium channel blockers, CsA is associated with gingival overgrowth with the wide range of prevalence rate estimated between 8% and 70%.² Despite the intensive studies investigating the cellular and molecular basis of the development of CsA-induced gingival overgrowth, the exact mechanism underlying this condition is still unclear. The imbalance of extracellular matrix molecules accumulation,^{3–9} upregulation of epithelial–mesenchymal transition (EMT) makers,^{10–12} and even the overexpression of embryonic stem cell (ESC) marker Nanog¹³ are believed to attribute the pathogenesis of CsA-induced gingival overgrowth.

Oct4, a key transcription factor, is involved in the maintenance of pluripotency and self-renewal in undifferentiated ESCs.^{14,15} Recently, Oct4 was found in fibroblasts derived from the cesarean scar tissue¹⁶ and keloid.¹⁷ In the present study, the effects of CsA on normal human gingival fibroblasts (HGFs) were determined through quantitative real-time reverse transcription-polymerase chain reaction (qRT-PCR) and western blot to determine the expression of Oct4. In addition, cell growth in CsA-treated HGFs with lentiviral-mediated shRNAi knockdown was evaluated by tetrazolium bromide reduction assay.

Materials and methods

Cell culture

Normal HGFs derived from three healthy male individuals undergoing surgical crown lengthening were cultured according to previous criteria and an explant technique as described previously.^{4,13} This study was approved by the Institutional Review Board permission at the Chung Shan Medical University Hospital.

Oct4 expression analyses

HGFs were arrested in G₀ by serum deprivation according to our previous experiments.^{3,5} Nearly confluent monolayers of HGFs were washed with serum-free medium and immediately exposed to 0, 100, 500, and 1000 ng/ml CsA (Sigma–Aldrich, St. Louis, MO, USA). Cell lysates were collected at 24 h for qRT-PCR and western blot assays. Cultures without fetal calf serum (FCS) (Gibco BRL, Gaithersburg, MD, USA) were used as negative control.

Quantitative real-time reverse transcription-polymerase chain reaction

Total RNA of cells was purified using Trizol reagent (Invitrogen Life Technologies, Carlsbad, CA, USA) according to our previous publication.^{10,11} Oct4 primers were designed: (forward) GTGGAGAGCAACTCCGATG and (reverse) TGCTC CAGCTTCTCCTTCTC. The glyceraldehydes-3-phosphate dehydrogenase (GAPDH) housekeeping gene was amplified

as a reference standard. *GAPDH* primers were designed: (forward) CATCATCCCTGCCTCTACTG and (reverse) GCCTGCTTACCACCTTC.

Western blot

The extraction of proteins from cells and immunoblotting procedure were performed as described previously.^{8,9} The primary antibodies were anti-Oct3/4 (SC-9081, Santa Cruz, CA, USA) and anti-GAPDH (#MA1-16757, ThermoFisher Scientific, Waltham, MA, USA) antibodies.

Oct4/Nanog knockdown in CsA-treated HGFs by lentiviral-mediated shRNAi

The pLV-RNAi vector was purchased from Biosettia Inc. (Biosettia, San Diego, CA, USA). The method of cloning the double-stranded shRNA sequence is described according to our recent publications.^{11,13} Lentiviral vectors expressing short hairpin RNA (shRNA) that targets human Oct4 (sh-Oct4-1: 5'-AAAAGCTGGGGAGAGTATATATTTTGGATCCAAAATATATACTCTCCCAGC-3'; sh-Oct4-2: 5'-AAAAGCTCTCCCATGCATTCAAA TTGGATCCAATTTGAATGCATGGGAGAGC -3'); Nanog (sh-Nanog: 5'-AAAAGCATCCGACTGTAAAGAATTTGGATCCAAATTCTTACAGTCGGATGC-3') were synthesized and cloned into pLV-RNAi to generate a lentiviral expression vector. shRNA that targets luciferase (sh-Luc: 5'-CCGGACTTACGCTGAGTACTTCCGAATCTCGAGTTCCGAAGTACTCAGCGTAAAGTTTTTTTG-3') was utilized for an experimental control.

Cell growth

HGFs placed in 96-well plates washed with phosphate-buffered saline and then cultured without FCS for starvation overnight. After treatment with 500 ng/ml CsA for 24 h, cell growth was tested using the MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay kit (Sigma–Aldrich, St. Louis, MO, USA) as described previously.¹³

Statistical analysis

All assays were repeated three times to ensure reproducibility. Statistical analysis was carried out by one-way analysis of variance (ANOVA). Tests of differences of the treatments were analyzed by Duncan's test. $P < 0.05$ was considered statistically significant.

Results

As shown in Fig. 1, CsA was found to increase Oct4 transcript in HGFs in a dose-dependent fashion ($p < 0.05$). CsA also upregulated the protein expression of Oct4 in a dose-dependent manner ($p < 0.05$) (Fig. 2). From the Alphascreen 2000, the amount of Oct4 was elevated about 1.2, 4.3, and 4.8 fold at concentrations of 100, 500, and 1000 ng/ml CsA, respectively, as compared with control.

Down-regulation of Oct4 in CsA-treated HGFs was achieved by viral transduction with lentiviral vector expressing shRNA targeting Oct4. In addition, lentiviral vector expressing sh-Luc was used as control. qRT-PCR confirmed




Figure 1 HGFs were treated with indicated concentration of CsA for 24 h. The Oct4 mRNA expression was examined by qRT-PCR. The relative Oct4 mRNA expression represent the mean \pm SD. * represents significant difference from control values with $p < 0.05$.

that lentivirus expressing both sh-Oct4 markedly reduced the expression level of CsA-induced Oct4 expression in HGFs ($p < 0.05$) (Fig. 3). As shown in Fig. 4, western blot confirmed that the knockdown of Oct4 could reduce Oct4 expression in CsA-stimulated HGFs. The quantitative measurement of Nanog protein expression by the Alphasizer 2000 is shown in the lower panel of Fig. 4.




Figure 2 The Oct4 protein expression was examined by western blot. HGFs were treated with indicated concentration of CsA for 24 h. GAPDH was used as protein loading control (Upper panel). Levels of Oct4 protein treated with CsA were measured by Alphasizer 2000. The relative level of Oct4 protein expression for each sample was normalized against GAPDH signal, and the control was set as 1.0. Triplicate experiments were performed. * represents significant difference from control values with $p < 0.05$ (Lower panel).




Figure 3 The silencing effect of Oct4 shRNA in CsA-treated HGFs was validated transcriptionally by qRT-PCR. Single cell suspension of HGFs was transduced with sh-Luc or sh-Oct4 lentivirus, individually or concurrently, and treated with or without CsA (500 ng/ml) as indicated. The relative Oct4 mRNA expression represent the mean \pm SD. * $p < 0.05$, Sh-Luc + CsA group vs. control group; # $p < 0.05$, Sh-Oct4-1 + CsA or Sh-Oct4-2 + CsA vs. Sh-Luc + CsA group.




Figure 4 Total proteins prepared from single cell suspension of HGF transduced with sh-Luc or sh-Oct4 lentivirus, individually or concurrently, and treated with or without CsA were analyzed for CsA expression by western blot. GAPDH was used as protein loading control. Levels of Oct4 protein from HGFs transduced with sh-Luc or sh-Oct4 lentivirus were measured by Alphasizer 2000. Triplicate experiments were performed. * represents significant difference from control values with $p < 0.05$. # $p < 0.05$, Sh-Oct4-1 + CsA or Sh-Oct4-2 + CsA vs. Sh-Luc + CsA group.

As shown in Fig. 5, the suppression of Oct4 was insufficient to prevent the CsA-induced cell growth. Our previous study¹³ has found that Nanog was associated with CsA-induced gingival overgrowth. Both Oct4 and Nanog are crucial pluripotency-associated transcription factors, we therefore utilized higher efficient sh-Oct4-2/sh-Nanog-1¹³ lentivirus and conducted the double knockdown experiments. The results demonstrated that the down-regulation of Oct4/Nanog significantly inhibited CsA-stimulated cell growth ($p < 0.05$).

Discussion

Gingival overgrowth is a widely reported side-effect of CsA therapy. Until now, the pathogenesis of CsA-induced gingival overgrowth still remains to be elucidated. Recently, *in situ* expression of OCT4, SOX2, KLF4, and C-MYC (OSKM) transcription factors during wound healing could diminish fibrotic activity and lead to reduce scar tissue formation in a mouse model.¹⁸ To the best of our knowledge, this is the first report showed that Oct4 mRNA and protein expression was increased after CsA treatment in HGFs. These findings may raise a question of whether Oct4 is involved in the enhanced cell proliferation following CsA administration.

Oct4 was reported to regulate tumor initiating property and EMT traits.¹⁹ EMT is critical for the development and the diseases including drug-induced gingival overgrowth.²⁰ Recently, the upregulation of Snail¹⁰ and Slug^{11,12} were found to play an important role in the pathogenesis of CsA-induced gingival overgrowth. Thus, the detailed molecular mechanisms involved in the regulatory links between Oct4 and EMT properties are worthy of further investigation.

Moreover, overexpression of Oct4 was found to enhance cell proliferative activity, invasiveness and colony formation in oral squamous cell carcinoma cell lines *in vitro*.¹⁹ In addition, Oct4 knockdown treatment could significantly slow down the tumor growth mediated in subcutaneous xenografts nude mice model.¹⁹ These studies indicated that Oct4 may participate in the regulation of oral cell growth and the inhibition of

Oct4 could attenuate their excessive growth. Hence, we utilized the lentivirus expressing sh-Oct4 to inhibit the levels of CsA-induced Oct4 transcript and protein expression in HGFs after CsA treatment and examine cell proliferation to assess the effect of Oct4 on CsA-induced gingival overgrowth. Surprisingly, we found that knockdown of Oct4 alone could not suppress CsA-stimulated HGFs growth. Our previous study has revealed that Nanog was increased in CsA-treated HGFs.¹³ Nanog and Oct4 have been reported to work in concert to support stem cell potency and self-renewal.¹⁵ Therefore, we conducted the double knockdown experiments with Oct4 and Nanog. The data revealed that co-knockdown with Oct4 and Nanog was able to attenuate the CsA-stimulated HGFs growth. Although the detailed mechanism regarding the interaction between these two factors requires further experiments to unveil, our results suggested that Oct4 may be just one of the downstream factors affected by CsA treatment during the progressive change of gingival overgrowth instead of the crucial initiator to control cell proliferation. However, further exploration of the mechanisms through which Oct4/Nanog regulates cell proliferation is still necessary to shed light on the role of Oct4/Nanog in the pathogenesis of CsA-induced gingival overgrowth.

In summary, this study demonstrated that the pluripotency-associated transcription factor Oct4 was elevated after CsA treatment, but this upregulation was not directly associated with excessive cell proliferation. It is reasonable to assume that the increase in Oct4 with CsA treatment may lead to the enhanced stemness in HGFs. Further studies are required to confirm this hypothesis. The interplay between Nanog and Oct4 in CsA-treated HGFs is worthy of further investigation.

Declaration of Competing Interest

None declared.

Acknowledgment

The work was supported by the National Science Council, Taiwan (NSC 101-2314-B-040-012).

References

1. Faulds D, Goa KL, Benfield P. Cyclosporin. A review of its pharmacodynamic and pharmacokinetic properties, and therapeutic use in immunoregulatory disorders. *Drugs* 1993;45: 953–1040.
2. Kataoka M, Kido J, Shinohara Y, Nagata T. Drug-induced gingival overgrowth-A review. *Biol Pharm Bull* 2005;28: 1817–21.
3. Lin HJ, Tsai CH, Huang FM, Chang YC. The upregulation of type I plasminogen activator inhibitor in human gingival fibroblasts stimulated with cyclosporin A. *J Periodont Res* 2007;42:39–44.
4. Tsai CH, Chang TY, Chang YC. Upregulation of lysyl oxidase expression in cyclosporin A-induced gingival overgrowth. *J Dent Sci* 2009;4:13–7.
5. Tsai CH, Yang SF, Huang FM, Chang YC. The upregulation of cystatin C in human gingival fibroblasts stimulated with cyclosporin A. *J Periodont Res* 2009;44:459–64.




Figure 5 The cell viability of the control or Oct4/Nanog-knockdown CsA-treated HGFs was evaluated by an MTT assay. Cells were exposed for 24 h in medium containing 500 ng/ml CsA. * $p < 0.05$, Sh-Luc + CsA group versus control group. # $p < 0.05$, Sh-Oct-4-2 + Sh-Nanog-1 + CsA vs. Sh-Luc + CsA group.

6. Ho YC, Lin HJ, Tsai CH, Chang YC. Regulation of type I plasminogen activator inhibitor in human gingival fibroblasts with cyclosporine A. *Oral Dis* 2010;16:396–401.
7. Chang TY, Tsai CH, Chang YC. The upregulation of heat shock protein 47 in human gingival fibroblasts stimulated with cyclosporin A. *J Periodont Res* 2010;45:317–22.
8. Lee SS, Tsai CH, Kuan YH, Huang FM, Chang YC. The upregulation of transglutaminase-2 by cyclosporine A in human gingival fibroblasts is augmented by oxidative stress. *J Periodontol* 2013;84:1469–75.
9. Tsai CH, Lee SS, Huang FM, Yu CC, Yang SF, Chang YC. The modulation of hypoxia-inducible factor-1 α /plasminogen activator inhibitor-1 axis in human gingival fibroblasts stimulated with cyclosporine A. *J Formos Med Assoc* 2015;114:58–63.
10. Lin YH, Yu CC, Lee SS, Chang YC. Elevated Snail expression in human gingival fibroblasts by cyclosporine A as the possible pathogenesis for gingival overgrowth. *J Formos Med Assoc* 2015;114:1181–6.
11. Tsai CH, Yu CC, Lee SS, Yu HC, Huang FM, Chang YC. Upregulation of Slug expression by cyclosporine A contributes to the pathogenesis of gingival overgrowth. *J Formos Med Assoc* 2016;115:602–8.
12. Lin T, Yu CC, Hsieh PL, Liao YW, Yu CH, Chen CJ. Down-regulation of miR-200b-targeting Slug axis by cyclosporine A in human gingival fibroblasts. *J Formos Med Assoc* 2018;117:1072–7.
13. Yu CC, Su NY, Liu CM, Yang LC, Tsai CH, Chang YC. The upregulation of embryonic stem cell marker Nanog in human gingival fibroblasts stimulated with cyclosporine A - an *in vitro* study. *J Dent Sci* 2017;12:78–82.
14. Jaenisch R, Young R. Stem cells, the molecular circuitry of pluripotency and nuclear reprogramming. *Cell* 2008;132:567–82.
15. Park IH, Zhao R, West JA, et al. Reprogramming of human somatic cells to pluripotency with defined factors. *Nature* 2008;451:141–6.
16. Rungsiwut R, Pavarajarn W, Numchaisrika P, Virutamasen P, Pruksananonda K. Transgene-free human induced pluripotent stem cell line (HS5-SV.hiPS) generated from cesarean scar-derived fibroblasts. *Stem Cell Res* 2016;16:10–3.
17. Zhang YX, Liu LP, Li M, et al. Development of individualized induced pluripotent stem cells from fibroblasts of keloid lesions in patients. *Transplant Proc* 2018;50:2868–71.
18. Doeser MC, Schöler HR, Wu G. Reduction of fibrosis and scar formation by partial reprogramming *in vivo*. *Stem Cells* 2018;36:1216–25.
19. Tsai LL, Hu FW, Lee SS, Yu CH, Yu CC, Chang YC. Oct4 mediates tumor initiating properties in oral squamous cell carcinomas through the regulation of epithelial-mesenchymal transition. *PLoS One* 2014;9:e87207.
20. Sume SS, Kantarci A, Lee A, Hasturk H, Trackman PC. Epithelial to mesenchymal transition in gingival overgrowth. *Am J Pathol* 2010;177:208–18.